



# A note on definition of matrix convex functions<sup>☆</sup>

Oleg E. Tikhonov

*Research Institute of Mathematics and Mechanics, Kazan State University, Nuzhina 17, Kazan 420008, Russia*

Received 6 May 2005; accepted 20 December 2005

Available online 8 February 2006

Submitted by C. Davis

## Abstract

We prove that a real-valued function  $f$  defined on an interval  $S$  in  $\mathbf{R}$  is matrix convex if and only if for any natural  $k$ , for all families of positive operators  $\{A_i\}_{i=1}^k$  in a finite-dimensional Hilbert space, such that  $\sum_{i=1}^k A_i = 1$ , and arbitrary numbers  $x_i \in S$ , the inequality

$$f\left(\sum_{i=1}^k x_i A_i\right) \leq \sum_{i=1}^k f(x_i) A_i$$

holds true.

© 2006 Elsevier Inc. All rights reserved.

**Keywords:** Matrix convex function; The Neumark theorem

The aim of this note is to add one more characterization to the list of known characterizations of matrix and operator convex functions (see, e. g., [1,3,4]).

In what follows,  $S$  stands for an interval in  $\mathbf{R}$  of an arbitrary type. For a function  $f: S \rightarrow \mathbf{R}$  and a self-adjoint operator  $X$  in a finite-dimensional Hilbert space  $H$  with spectrum in  $S$ , the value  $f(X)$  is defined by the spectral theorem. The identity operator in  $H$  is denoted by  $1_H$ .

**Theorem.** *For a function  $f: S \rightarrow \mathbf{R}$ , the following conditions are equivalent:*

(i)  $f$  is matrix convex, i.e.,

$$f(\alpha X + (1 - \alpha)Y) \leq \alpha f(X) + (1 - \alpha)f(Y)$$

<sup>☆</sup> Supported by the Russian Foundation for Basic Research (grant no. 05-01-00799).

E-mail address: [Oleg.Tikhonov@ksu.ru](mailto:Oleg.Tikhonov@ksu.ru)

for all self-adjoint operators  $X, Y$  in a finite-dimensional Hilbert space and every  $\alpha \in [0, 1]$ , provided that spectra of  $X$  and  $Y$  lie in  $S$ ;

- (ii) for any natural  $k$ , for all families of positive operators  $\{A_i\}_{i=1}^k$  in a finite-dimensional Hilbert space  $H$ , such that  $\sum_{i=1}^k A_i = 1_H$ , and arbitrary numbers  $x_i \in S$ , the inequality

$$f\left(\sum_{i=1}^k x_i A_i\right) \leq \sum_{i=1}^k f(x_i) A_i$$

holds true.

**Proof.** Since

$$(\min_i x_i) 1_H \leq \sum_{i=1}^k x_i A_i \leq (\max_i x_i) 1_H,$$

the left hand side in the inequality in (ii) is well-defined.

(ii)  $\implies$  (i). Let  $H$  denote the space where the operators  $X$  and  $Y$  act,

$$X = \sum_{i \in I} \lambda_i P_i \quad \text{and} \quad Y = \sum_{j \in J} \mu_j Q_j$$

be their spectral decompositions. Then

$$\sum_{i \in I} \alpha P_i + \sum_{j \in J} (1 - \alpha) Q_j = 1_H$$

and

$$\begin{aligned} f(\alpha X + (1 - \alpha)Y) &= f\left(\sum_{i \in I} \lambda_i \alpha P_i + \sum_{j \in J} \mu_j (1 - \alpha) Q_j\right) \\ &\leq \sum_{i \in I} f(\lambda_i) \alpha P_i + \sum_{j \in J} f(\mu_j) (1 - \alpha) Q_j \\ &= \alpha f(X) + (1 - \alpha) f(Y). \end{aligned}$$

(i)  $\implies$  (ii). It is easy to see that without loss of generality we can suppose that  $0 \in S$ . By the Neumark theorem [5], there exists a space  $\mathcal{H}$  larger than  $H$  and a family of mutually orthogonal projections  $P_i$  in  $\mathcal{H}$  such that  $\sum_{i=1}^k P_i = 1_{\mathcal{H}}$  and  $A_i = P P_i P|_H$  ( $i = 1, \dots, k$ ), where  $P$  is the projection from  $\mathcal{H}$  onto  $H$ . Then we have

$$\begin{aligned} f\left(\sum_{i=1}^k x_i A_i\right) &= f\left(\sum_{i=1}^k x_i P P_i P|_H\right) = f\left(P\left(\sum_{i=1}^k x_i P_i\right) P|_H\right) \\ &= P f\left(P\left(\sum_{i=1}^k x_i P_i\right) P\right) P|_H \\ &\leq P f\left(\sum_{i=1}^k x_i P_i\right) P|_H = P\left(\sum_{i=1}^k f(x_i) P_i\right) P|_H \\ &= \sum_{i=1}^k f(x_i) A_i. \quad \square \end{aligned}$$

(The inequality in the above calculation follows from the Davis inequality [2]:  $Pf(PXP)P \leq Pf(X)P$ .)

**Remark.** After the first version of the present paper had been submitted, professor T. Ando recommended to add one more equivalent condition:

(iii) for any natural  $k$ , for any family of operators  $\{B_i\}_{i=1}^k$  with spectra in  $S$  and for any family of operators  $\{A_i\}_{i=1}^k$  such that  $\sum_{i=1}^k A_i^* A_i = 1_H$  the inequality

$$f\left(\sum_{i=1}^k A_i^* B_i A_i\right) \leq \sum_{i=1}^k A_i^* f(B_i) A_i$$

holds.

He gave a proof of the implications (ii)  $\iff$  (iii) (one can read that proof in [6]). Since the equivalence of (i) and (iii) is known (it goes back to [3]), the proof due to Ando gives an alternative proof of the theorem presented in this note. It is worth to note that in contrast to (i)  $\implies$  (ii) and (i)  $\implies$  (iii), the proof of (ii)  $\iff$  (iii) does not refer to the Neumark theorem and deals with one and the same finite-dimensional Hilbert space.

## Acknowledgments

The author thanks T. Ando for a useful comment on the original version of the paper.

## References

- [1] J. Benda, S. Sherman, Monotone and convex operator functions, *Trans. Amer. Math. Soc.* 79 (1955) 58–71.
- [2] C. Davis, A Schwarz inequality for convex operator functions, *Proc. Amer. Math. Soc.* 8 (1957) 42–44.
- [3] C. Davis, Notions generalizing convexity for functions defined on spaces of matrices, *Proc. Amer. Math. Soc. Symposia (Convexity)* 7 (1963) 187–201.
- [4] F. Hansen, G.K. Pedersen, Jensen's inequality for operators and Löwner's theorem, *Math. Ann.* 258 (1982) 229–241.
- [5] M.A. Neumark, On a representation of additive operator set functions, *Dokl. Akad. Nauk SSSR* 41 (9) (1943) 373–375. (Russian); English translation: C.R. (Doklady) Akad. Sci. USSR. (N.S.) 41 (1943), 359–361.
- [6] O.E. Tikhonov, A note on definition of matrix convex functions with a supplement by T. Ando, Preprints of the N.G. Chebotarev Research Institute of Mathematics and Mechanics of Kazan State University, Preprint 0003-2005. Available from: <<http://www.niimm.ksu.ru/data/preprints/thepreprints/0003-2005.ps>>.